
Test contains 15 questions, 2 marks each. **No negative marks.**

$N_A = 6 \times 10^{23} \text{ mol}^{-1}$; Ar: N-14; O-16; K-39; I-127.

1. Electron configuration for the element E is $1s^2 2s^2 2p^6 3s^2 3p^6$.
Which one of the following statements is correct?
 - 1) Mass number of the element E is 18
 - 2) Atom of the element E contains 6 valence electrons
 - 3) The element E is in Group VIA and Period 3
 - 4) Valence electrons of the element E are in the third shell

2. Calculate the value of the equilibrium constant, K_c , for the reaction:
$$2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$$

for given equilibrium concentration: HI 0.1 mol/L, H_2 0.4 mol/L and I_2 0.2 mol/L.
 - 1) 8
 - 2) 0.125
 - 3) 1.25 L/mol
 - 4) 0.8 mol/L

3. Calculate the mass percent (%) of NaOH in a solution prepared by adding 100 g of water to 200 g 30 % NaOH solution.
 - 1) 15 %
 - 2) 10 %
 - 3) 20 %
 - 4) 25 %

4. In oxidation-reduction reaction between potassium iodide and potassium nitrite in acidic solution (H_2SO_4) produce elemental iodine, nitrogen monoxide, potassium sulfate and water.
If 25 mL 2 mol/L of potassium nitrite solution is reacted, how many grams of iodine are produced?
 - 1) 12.7
 - 2) 6.35
 - 3) 3.175
 - 4) 25.4

5. Which of the following water solution is basic?

- 1) Solution which in 0.1 L contains 10^{-8} mol OH^-
- 2) Solution which in 10 mL contains 10^{-8} mol H^+
- 3) Solution which in 1 L contains 6×10^{18} OH^-
- 4) Solution with a $\text{pH}=3$

6. In which one of the following sets all substances have pH of water solutions higher than pH of pure water?

- 1) CaO , Na , NaCH_3COO
- 2) Na_2CO_3 , NaNO_3 , CO_2
- 3) SO_2 , NH_4Cl , CH_3COOH
- 4) NaNO_2 , HNO_2 , N_2O_3

7. Which of the following substances reacts with hydrochloride acid?

- 1) NH_4Cl
- 2) NaCH_3COO
- 3) elemental silver
- 4) CO_2

8. Mark **the correct** statement:

- 1) benzene has three single (C-C) longer bonds and three double (C=C) shorter bonds
- 2) length of all bonds in benzene is between the length of the single (C-C) bond and the length of the double (C=C) bond
- 3) benzene reacts with hydrochloric acid
- 4) reaction of benzene and an electrophile produces carbanion in the initial step

9. Mark **the correct** statement:

- 1) addition of sulfuric acid to alkenes follows the ionic mechanism
- 2) addition of sulfuric acid to alkenes is initiated by formation of a carbanion
- 3) addition of sulfuric acid to alkenes is initiated by nucleophilic addition of the sulfate anion
- 4) addition of sulfuric acid to alkenes does not bear any similarities to the addition of hydrochloric acid to alkenes

10. In which group **all listed compounds** react with hydrogen in the presence of a catalyst:

- 1) cyclobutane, propene, cyclopentane
- 2) cyclopropane, isobutane, 1-butanol
- 3) cyclopentane, cyclopropane, cyclohexene
- 4) cyclopentene, cyclopropane, propanone

11. Thermal cracking of alkanes can be considered as:

- 1) elimination reaction
- 2) substitution reaction
- 3) acid-base reaction
- 4) reduction with hydrogen

12. Mark the **correct** statement:

- 1) aniline is stronger base than ammonia
- 2) pyridine is less reactive than benzene in electrophilic aromatic substitution reactions
- 3) amides form stable salts in reaction with acids
- 4) pyrrole has properties of secondary amines

13. What is molecular formula of the product obtained in reaction of a single molecule of methanol and a single molecule of phosphoric acid:

- 1) $\text{CH}_5\text{O}_4\text{P}$
- 2) $\text{C}_3\text{H}_9\text{O}_4\text{P}$
- 3) $\text{CH}_7\text{O}_5\text{P}$
- 4) $\text{CH}_4\text{O}_3\text{P}$

14. Reduction of pyridine by H_2 in the presence of catalyst affords:

- 1) pyrrolidine
- 2) tertiary amine
- 3) the product more basic than pyridine
- 4) purine derivative

15. Mark the **incorrect** statement related to uronic acids:

- 1) in solution they may form β -anomer in pyranose form
- 2) in reaction with alcohols they form acetals
- 3) they do not show reducing properties
- 4) they are formed by oxidation of the primary alcoholic group of aldoses